

Estimation of the Latitude, the Gnomon's Length and Position About *Sinbeop-Jipyong-Ilgu* in the Late of Joseon Dynasty

Byeong-Hee Mihn^{1,2}, Yong Sam Lee^{3†}, Sang Hyuk Kim^{1,2}, Won-Ho Choi⁴, Seon Young Ham^{1,3}

¹Korea Astronomy and Space Science Institute, Daejeon 34055, Korea

²Korea University of Science and Technology, Daejeon 34113, Korea

³Department of Astronomy and Space Science, Chungbuk National University, Cheongju 28644, Korea

⁴The Research Institute of Digital Heritage, Jeonju University, Jeonju 55069, Korea

In this study, the characteristics of a horizontal sundial from the Joseon Dynasty were investigated. Korea's Treasure No. 840 (T840) is a Western-style horizontal sundial where hour-lines and solar-term-lines are engraved. The inscription of this sundial indicates that the latitude (altitude of the north celestial pole) is $37^{\circ} 39'$, but the gnomon is lost. In the present study, the latitude of the sundial and the length of the gnomon were estimated based only on the hour-lines and solar-term-lines of the horizontal sundial. When statistically calculated from the convergent point obtained by extending the hour-lines, the latitude of this sundial was $37^{\circ} 15' \pm 26'$, which showed a $24'$ difference from the record of the inscription. When it was also assumed that a convergent point is changeable, the estimation of the sundial's latitude was found to be sensitive to the variation of this point. This study found that T840 used a vertical gnomon, that is, perpendicular to the horizontal plane, rather than an inclined triangular gnomon, and a horn-shaped mark like a vertical gnomon is cut on its surface. The length of the gnomon engraved on the artifact was 43.1 mm, and in the present study was statistically calculated as 43.7 ± 0.7 mm. In addition, the position of the gnomon according to the original inscription and our calculation showed an error of 0.3 mm.

Keywords: horizontal sundial (*jipyong-ilgu*), gnomon, hour-line, solar-term-line, sundial's latitude

1. INTRODUCTION

From the 17th century, Joseon began to import and employ Western science practices from China (Yu 1983; Lee 1985; Hahn 2004). The *Shixian* Calendar was introduced and established in Joseon in 1654 and this marked the beginning of the practice of Western science in the field of Joseon's royal politics (Yu 1982; Lee et al. 2010). The continuous influx of Western science also had an influence on the astronomical observation instruments utilized in the late Joseon Dynasty. A representative example is the manufacture of a Western-style horizontal sundial (新法地平日晷) (Hahn 2004).

The sundials of the Joseon Dynasty could measure the time and 24 solar terms together. The various sundials developed in the reign of King Sejong were mostly equatorial sundials

(Lee & Kim 2011; Kim et al. 2012; Lee et al. 2016). However, *Angbuilgu*, a scaphe sundial was installed within Hanyang, the capital of this kingdom (Seoul), and thus was a sundial that could be used by the public as well as the palace dwellers. This scaphe was the modification of *Yangyi* (仰儀), which is a large-scale observation instrument of the Yuan (元) Dynasty. Interestingly, hour-lines and solar-term-lines are drawn on this hemispherical sundial. Thanks to *Angbuilgu*, it was recognized that a sundial indicates the time of a day and the date (or season) of a year in Joseon. This perception seems to have motivated Crown Prince Sohyeon (1612~1645) to introduce sundials to his country when he received a Western-style horizontal sundial as a gift from Adam Schall von Bell around 1644 (Lee 1985).

The hour-lines and solar-term-lines of Western-style hori-

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received 19 APR 2017 Revised 5 JUN 2017 Accepted 7 JUN 2017

†Corresponding Author

Tel: +82-43-261-2312, E-mail: leeysam@hanmail.net

ORCID: <https://orcid.org/0000-0002-4878-8664>

zontal sundials featured a new format which had never been used in the early *Joseon* Dynasty. It seems that a Western-style horizontal sundial imported by Crown Prince Sohyeon could not be immediately reconstructed into another version in accordance with Hanyang. According to the record of *Gukjo-ryeoksang-go* (國朝曆象考, Reference for Astronomy and its Calendar of Our Country), Kim Young (1749~1817) made a horizontal sundial in 1789. Also, it explains the hour-lines and solar-term-lines of this sundial in detail (Lee & Moon 2004). Besides Chen & Hua (2011) has described a vertical gnomon related to a drawing of a horizontal sundial with an old Chinese method which is similar to a drawing method of the record of *Gukjo-ryeoksang-go*.

In Korea, some extant Western-style horizontal sundials have been designated as treasures. A black stone horizontal sundial and a white marble horizontal sundial were once displayed in the King Sejong Great Memorial Hall (Lee 1984). They have been designated as Treasure No. 840 (hereafter 'T840') and Treasure No. 839 (hereafter 'T839') respectively, and are currently housed in the national palace museum of Korea (NPM). The former has an inscription that records the latitude (the altitude of the north celestial pole) of Hanyang, and the latter has an inscription that is thought to relate details about the manufacturing date and the sundial maker. For these stone horizontal sundials, the gnomon is lost, and thus its shape is not known. In addition, an artifact made of copper is owned by the Seoul national university museum (SNUM) (Song et al. 1994). This artifact could be related to the metallic horizontal sundial that was manufactured by Kim Young as mentioned in *Gukjo-ryeoksang-go*.

Lee (1984) investigated the external characteristics of the two stone horizontal sundials. Lee (1985) also discussed the permanent residence of the white marble horizontal sundial (T839), and calculated the latitude of this sundial with the

rubber of this horizontal sundial. We received the 3D scan data of T840 from the cultural heritage administration of Korea (CHA). In the present study, the latitude of the black stone horizontal sundial (T840) and the length of the stylus were analyzed using the 3D scan data with a point density of 1 mm. In Chapter 2, the 3D scan data and the analytic method are discussed. In Chapters 3 and 4, the latitude and stylus of the T840 horizontal sundial are analyzed and compared based on the inscription engraved on the original artifact.

2. DATA REDUCTION AND CALCULATION

2.1 Three-dimensional Scan Data

To obtain detailed information on the lines and features of the horizontal sundial, a three-dimensional scanner was used with the optical triangulation method. The optical triangulation method can be classified into a fringe projection type and an optical triangulation laser type. The former obtains the information through illuminating a light source onto the object, and the latter acquires the information on an object by shooting a laser (Choi & Ko 2017).

In the case of the T840 horizontal sundial, 3D scan measures were conducted in 2015 using the ATOS Compact Scan 5M (GOM, Germany). This is based on the fringe projection-type phase shift pattern method. The data precision of the 3D scanner was 0.001 mm ~ 0.02 mm, and the resolution of the camera was 2,448 × 2,050 pixels. The projection method was the phase shift pattern method, and the light source was blue light-emitting diode (LED). Based on the aforementioned settings, high-resolution 3D data could be obtained. The fringe projection-type 3D scanner (Fig. 1(a)) consisted of a light source, an illuminant lens, and a charge coupled device (CCD) (Lee et al. 2012).




Fig. 1. (a) The diagram of a fringe projection (Lee et al. 2012), (b) the image detection and processing of a 3D scan.

In the fringe projection measurement, a dispersive light through the lens illuminates an object, and the CCD camera detects the illuminated light reflected from the object. At the same time, the depth pattern of an object surface is analyzed from the detected light, and the three-dimensional shape of the object surface is digitized. Fig. 1(b) shows the 3D scanning and analysis process for the horizontal sundial used in the present study. The work process consists of the 3D scanning stage and modeling stage for the horizontal sundial. In the scanning stage, the measurement area of the object is determined at first and then scanned at the optimum distance of the scanner for that area (Choi & Ko 2017). The raw scan data is converted into an editable data format. The data is optimized to 3D points through a noise removing process and a hole filling process. The 3D point data of each detection area is optimized and merged to finally become polygon data (Choi & Ko 2017). Through these modeling stages the polygon data contains the depth information of the sundial's surface where the hour-lines and solar-term curves are engraved. For the colors according to the depth ranges,




Fig. 2. Treasure No. 840's complanation image of 3D scan data.

the deeper the part is, the darker it gets, and the shallower the part is, the brighter it gets. When the polygon image is assigned black intensity depending on the depth, it becomes a complanation image such as the one shown in Fig. 2.

Fig. 2 shows the complanation image of T840 (hereafter 'T840-plate'). As in the original form of the T840-plate, the center line of all hour-lines (i.e., 12^h-line) are rotated by about 17.242° in the clockwise direction relative to the vertical line of the black stone. This sundial has a total of 13 hour-lines (every hour from 6:00 to 18:00), and a total of 11 solar-term-lines that represent the 24 solar terms. However, there are not 13 solar-term-lines. A curve for Bearded grain (芒種) and Great heat (大暑) are used as the summer solstice line together, and in the same manner a curve for Heavy snow (大雪) and Slight cold (小寒) are used as the winter solstice line. In addition, there is a horn-shaped groove with a length of 43.1 mm below the inscription that reads '新法地平日晷' indicating a Western-style horizontal sundial. An inscription that states that the latitude of this sundial is 37° 39' is engraved on the southern lateral face of this horizontal sundial (i.e., the upper side of Fig. 2).

2.2 Analytical Methods

The coordinate system of the horizontal sundial can be conveniently specified as shown in Fig. 2. The origin of this coordinate system is identical to the convergent point obtained by extending the hour-lines of a horizontal sundial (Lee 1982). It is generally set up so that the west in the horizontal plane is determined as the X-axis, and the south as the Y-axis. Fig. 3 shows the diagram for the gnomon's projection for a horizontal sundial. Point A is the position of the gnomon, and point C is the position where the shadow of the gnomon's tip is cast, that lies on a line of the vernal




Fig. 3. (a) The coordinate of a horizontal sundial and (b) diagram for a gnomon's projection by the sun.

and autumnal equinoxes. The direction from the origin toward the gnomon's tip is the North celestial pole, and thus the angle between \overline{OB} and a horizon is identical to the latitude of its sundial. When the length of the gnomon is H and the declination of the sun is δ , D is a shadow position of the gnomon's tip on the 12^h-line and \overline{AD} is N .

Fig. 4 shows the convergent point of each hour-line in the T840-plate. When the convergent point was magnified, it is not only one point, but is a lot of intersection points. Owing to an eccentric error of hour lines, the intersection points between the noon line (12^h-line) and the other hour-lines are dispersed from a virtual convergent point. In Fig. 4(b), each intersection point was expressed as symbols (P1, P2, P3, ..., M1, M2, M3..., P6 = M6). The center point of the east-west line (AA) is also one of the intersection points. If these intersection points are eligible to be a convergent point, then they are assumed as the origins of the sundial coordinate system. These origins have characteristics similar to the convergent point's samples which can be analyzed statistically. In other words, when each intersection point lies as the origin of the sundial coordinate, the latitude of the sundial and the length of the gnomon can be inversely estimated from the hour-lines and solar-term-lines at each origin.

The inscription of T840 specifies the latitude of Hanyang (Seoul) installed in this sundial. For each origin shown in Fig. 4(b), the latitude of the sundial can be estimated based on the angle between the 12^h-line and the hour-line (z) shown in Fig. 4(a) (Lee 1982; Rohr 1996).

$$\sin \phi_j = \frac{\tan z_{ij}}{\tan HA_j} \quad (i = o_1, o_2, o_3, \dots, j = 1, 2, 3, \dots) \quad (1)$$

where ϕ is the calculated latitude of the sundial, HA is the hour angle, and z is the angle between the 12^h-line and the

hour-line. The dummy index i represents the 12 origins, and j represents the 12 hour angles. In Eq. (1), the mean latitude of the sundial for each origin can be calculated as $\phi_i = \sum_j \phi_{ij}$.

On the other hand, the intersection point between the 12^h-line and a solar-term-line ($0, y$) corresponds to a point D_j ($j = 1, 2, 3, \dots$) in Fig. 3(b). The latitude of the sundial can also be calculated from two solar-term-lines with the same declination of the sun (Lee 1985).

$$\tan \phi_{ik} = \frac{y_{i,n+1-k} - y_{i,k}}{y_{i,n+1-k} + y_{i,k}} \cot \delta_{ik} \quad (i = o_1, o_2, o_3, \dots, k = 1, 2, 3, \dots) \quad (2)$$

where y_{ik} is the y value of each solar-term point on the 12^h-line measured from T840-plate. The dummy index i represents the 12 origins, and k represents the 6 solar-term-lines (declination of the sun).

Based on the mean latitude of the sundial obtained by Eq. (1) and the intersection points (x, y) of each solar-term-line and the hour-line, the length of the gnomon (H) can be calculated as follows.

$$H_{ijk} = (\cos \phi_i \cos HA_j + \sin \phi_i \tan \delta_{ik}) \sqrt{y_{ijk}^2 \sin^2 \phi_i - x_{ijk}^2} \quad (i = o_1, o_2, o_3, \dots, j = 1, 2, 3, \dots, k = 1, 2, 3, \dots) \quad (3)$$

where the dummy index i represents the 12 origins, j represents the 12 hour angles, and k represents the 6 solar-term-lines. In Eq. (3), the mean gnomon length for each solar-term-line ($H_{ik} = \sum_j H_{ijk}$) or the mean gnomon length for each hour-line ($H_{ij} = \sum_k H_{ijk}$) can be calculated for each origin. Also, the mean gnomon length for all origins is $H_i = \sum_{j,k} H_{ijk}$.

With the y_{ik} in Eq. (2) and the values in Eq. (3), the distance from each D_k on the 12^h-line to the gnomon (A), $N_k = \overline{AD}_k$ can be calculated as follows.

$$N_{ik} = \frac{H_{ik}}{\tan \phi_i} - y_{ik} \quad (i = o_1, o_2, o_3, \dots, k = 1, 2, 3, \dots) \quad (4)$$




Fig. 4. (a) A convergent point of all hour-lines in T840 and (b) an enlarged image of (a) at a convergent point.

The mean distance from each solar-term point on the 12^h-line to the gnomon can be calculated as $N_k = \sum_i N_{ik}$. In Fig. 3(b), $\overline{O_i A} = \overline{O_i D_k} - N_k$.

3. RESULTS

3.1 Latitude of the Sundial

The ϕ_{ij} calculated by Eq. (1) is analyzed statistically for a given origin. As shown in Fig. 5, the ϕ_{ij} for the origin AA could be indirectly compared with the actual trajectory of the mean latitude (ϕ_{AA}). The abscissa is the hour angle, and the ordinate is z . For each of the given origins of i , the ϕ_{ij} calculated for each hour angle was different from the mean latitude (ϕ_i), and they were consistent within a maximum of 42'. In this regard, the difference between z_{mean} and z_{ij} was several 10', where z_{mean} is an hour-line's angle calculated by mean latitude ϕ_{ij} for a j 's hour-line and the z_{ij} is j 's hour-line's angle measured from T840-plate. Regarding the origin of other i -s, the results were also similar to that shown in Fig. 5.

For every origin of Fig. 4(b), the mean latitude and $z_{ij} - z_{mean}$ of a j 's hour angle is shown in Fig. 6. The mean of each origin's mean latitude is $37^\circ 15' \pm 26'$, and this value embraces the latitude of T840 ($37^\circ 39'$). However, this mean for all origins showed a difference of about 24' compared to the actual value of the inscription.

3.2 Length of the Gnomon

The length of the gnomon for each origin can be estimated

by Eq. (3). This value is calculated using two variables: hour angle and the declination of the sun. Accordingly, for the statistical results, the length of the gnomon (H) can also be analyzed for each hour-line and for each solar-term-line, respectively. Let H_i be the mean gnomon length of i 's origin. Let H_{ij} be the mean gnomon length of i 's origin and j 's hour-line. Let H_{ik} also be the mean gnomon length of i 's origin and k 's solar-term-line. When the difference between H_i and H_{ij} was defined as ΔH_{ij} ($= H_{ij} - H_i$), the ΔH_{ij} is presented in Fig. 7. When the difference between H_i and H_{ik} is defined as ΔH_{ik} ($= H_{ik} - H_i$), the ΔH_{ik} is presented in Fig. 8. The abscissa is the hour angle, and the ordinate is the ΔH_{ij} in Fig. 7. The abscissa is the solar-term-line, and the ordinate is the H_{ik} in Fig. 8. The acronym of 24 solar terms is derived from the designation of them by Kim et al. (2017).

In Figs. 7 and 8, the ΔH_{ij} and ΔH_k showed slightly different distribution patterns relative to the H_i . In general, H_{ij} is more distributed than H_k relative to H_i . On the other hand, the standard deviation of each H_{ij} was smaller than that of each H_k . For example, in the case of Point AA, the mean of ΔH_{ij} was 0.3 mm, and that of ΔH_k was 0.2 mm. On the other hand, the mean of the standard deviations of H_{ij} was 0.4 mm, and that of H_k was 0.6 mm. The mean of the gnomon lengths by Eq. (3) for i 's origin (H_i) was 43.70 ± 0.66 mm. On T840-plate (Fig. 2), the horn-shaped groove (or pin mark) below the title's inscription reads 43.1 mm long. Although there is an error up to approximately 1 mm, the horn-shaped groove represents the length of the gnomon. It is possible that the gnomon of T840 sundial used a vertical stylus as shown in Fig. 3(b), rather than a triangular gnomon whose side is inclined from the horizontal plane by as much as the latitude of sundial (Lee 1985).




Fig. 5. The calculated latitudes of the sundial, T840, every hour angle about the origin AA.




Fig. 6. The differences between calculated mean z_{mean} and each measured hour-line angle z_j of T840-plate.




Fig. 7. $\Delta H_{ik}(= H_j - H_i)$ according to 13 hour-lines each origin.

3.3 Position of the Gnomon

T840-plate only has hour-lines and solar-term-lines without a gnomon. Each solar-term point on the 12^h-line can estimate the position of the gnomon using Eq. (4). Let N_k be the mean distance from the gnomon to each solar-term point on the

12^h-line. The distance from the winter solstice point to each solar-term point on the 12^h-line ($\overline{D_k D_{11}}$) can be measured from T840-plate. It can be compared with $N_k + \overline{D_i D_{11}}$ as summarized in Table 1. The mean value of $N_k + \overline{D_i D_{11}}$ is 76.29 ± 0.44 mm.

The $\overline{AD_{11}}$ measured from T840-plate was 76.0 mm. Therefore, the statistical position of the gnomon obtained by Eq. (4)




Fig. 8. $\Delta H_k (= H_k - H_i)$ according to 11 season-lines each origin.

Table 1. Comparison of gnomon's position

Name of season-line	N_k	$\overline{D_k D_{11}}$	$\overline{N_k + D_k D_{11}}$
Summer solstice (夏至) line	10.61 ± 0.23	66.08	76.69 ± 0.23
Grain fills (小滿) to Great heat (大暑) line	13.40 ± 0.22	62.80	76.20 ± 0.22
Summer begins (立夏) to Autumn begins (立秋) line	16.65 ± 0.20	59.55	76.20 ± 0.20
Crop rains (穀雨) to End of heat (處暑) line	21.15 ± 0.19	54.87	76.02 ± 0.19
Clear and bright (清明) to White dew (白露) line	26.78 ± 0.19	48.93	75.71 ± 0.19
Vernal equinox (春分) to Autumn equinox (秋分) line	32.77 ± 0.17	44.10	76.87 ± 0.17
Inserts awake (驚蟄) to Cold dew (寒露) line	41.01 ± 0.20	35.02	76.03 ± 0.20
Rainy weather (雨水) to Frost descends (霜降) line	49.31 ± 0.21	27.26	76.57 ± 0.21
Spring opens (立春) to Winter begins (立冬) line	59.48 ± 0.26	16.37	75.86 ± 0.26
Severe cold (大寒) to Slight snow (小雪) line	68.21 ± 0.30	7.99	76.20 ± 0.30
Winter solstice line	76.84 ± 0.34	0.00	76.84 ± 0.34

is consistent with the actual position of the gnomon on the T840 sundial.

4. DISCUSSIONS

As shown in Fig. 4(b), various origins were generated during the acquisition of the convergent point. For T840-plate, the other origins except AA were distributed on the extended line of the 12^h-line. If any origin is farther away from or close to the solar-term-lines, the statistical value may change to larger or smaller. The origin P1 is the closest to solar-term-lines, and the origin P3 is the farthest from them.

The mean latitude calculated is sensitive to changes in the origin. Fig. 9 shows the distribution of ϕ_i for each origin. The abscissa is the position of the origin, and the ordinate is

the mean latitude (ϕ_i) for each origin. The closer the origin is to the solar-term-lines, the larger the mean latitude is estimated by Eq. (1). The rate of the mean latitude change was -19.32°/mm. When two origins had a difference of 3 mm, the latitude between the two shows a difference of 1° statistically.

On the other hand, the ϕ_i obtained by Eq. (2) was 36° 55' ± 24'. This value was smaller than the mean latitude obtained by Eq. (1) (37° 15' ± 27'), and showed a larger difference from the real latitude of T840 (37° 39'). The rate of ϕ_i 's change by Eq. (2) was -18°/mm.

The mean length of the gnomon was impervious to the change in the origin. Fig. 10 shows the distribution of H_i for each origin. The farther the origin is from the solar-term-lines, the larger the mean length of the gnomon is estimated by Eq. (3). The rate of H_i 's change per 1.0 mm in the origin




Fig. 9. ϕ_i 's variation on the origin's offset based on the point AA.




Fig. 10. H_i 's variation on the origin's offset based on the point AA.

was 0.27 mm. When two origins had a difference of 3.6 mm, the length of the gnomon between the two statistically shows a difference of 1 mm.

Finally the position of the gnomon was also insensitive to the change in the origin. Fig. 11 shows the distribution of \overline{OA} for each origin (see Fig. 3). The farther the origin is from the solar-term-lines, the larger the \overline{OA} value is calculated by Eq. (4). The rate of \overline{OA} 's change per 1.0 mm in the origin was 1.12 mm.

5. CONCLUSIONS

T840 is a horizontal sundial that was newly manufactured considering the latitude of Hanyang (Lee 1984). This Western-style sundial with a missing gnomon is owned by the National Palace Museum of Korea. According to the studies of Lee (1982, 1985), the gnomon of this sundial is widely accepted as being triangular in shape, and the triangular gnomon is applied to its reconstruction in many museums.

In the present study, the latitude of T840 sundial is estimated using its 3D scan data provided by the Cultural Heritage




Fig. 11. \overline{OA} 's variation on the origin's offset based on the point AA.

Administration of Korea. By extending the hour-lines of T840-plate, the various origins are produced around the ideal convergent point. This study can verify whether the statistical results of a horizontal sundial are consistent with the inscription of T840 in which the latitude of Hanyang is specified as $37^{\circ} 39'$. This study can be summarized according to the following three main points.

First, the latitude of T840 was statistically estimated to be $37^{\circ} 15' \pm 26'$, which is consistent with the actual value in its inscription within a difference of about $24'$. However, the estimation of the latitude is very sensitive to variation of an origin, and the rate of its change was $-19.32'/\text{mm}$. For the latitude estimation, its calculation could be performed using only the solar-term point on the 12^{h} -line, but it showed a larger difference from the real value.

Second, the length and position of a vertical gnomon were consistent with the measure of T840-plate. It is generally understood that it is convenient to use a triangular gnomon to read a time from hour-lines on a horizontal sundial. However, if this sundial used to a vertical gnomon similar to Chen & Hua (2011), it is very interesting the fact that the length of the gnomon was consistent with the size of the cone-shaped groove engraved on the artifact. The length of the triangular cone-shaped groove was 43.1 mm, and our statistical result is 43.70 ± 0.66 mm. Also, the position of the gnomon showed an error of 0.29 mm. In addition, when the origin is varied by 1 mm, estimates of the length and position of the gnomon vary only 0.27 mm and 1.12 mm, respectively.

Third, this study found that T840 used a vertical stylus, that is, perpendicular to the horizontal plane, rather than an inclined triangular gnomon. This is why a triangular

gnomon needs to have at least two hole traces for the gnomon's installation where the two holes are placed on the extension of the 12^{h} -line. However, T840-plate has the trace of installation only at the position where the stylus is installed vertically.

We hope that this study will help in the investigation of the astronomical characteristics of a horizontal sundial which is remained only both straight lines and curved lines like T839. In the last decades in Korea, ones have made a lot of T840's replication which is measured by a triangular gnomon, not a vertical gnomon. It will also need to investigate more evidence why replicas of T840 were manufactured with a triangular gnomon, except the suggestion of Lee (1985).

ACKNOWLEDGMENTS

This work was supported by the research grant of the Chungbuk National University in 2014. We are especially grateful to the Cultural Heritage Administration of the Korea for providing 3-dimensional scan data of the extant Western-style horizontal sundials.

REFERENCES

- Chen MD, Hua TX, The Whole History of Chinese Astronomical Timekeeping Instrument (Ancient Volume) (Anhui Education Publishing, Hefei, 2011), 361-366.
- Choi WH, Ko SW, The epigraph reading method using a visualization technique based on morphological characteris-

- tics of the letters, J. Korea Contents Assoc. 17, 740-749 (2017).
- Hahn YH, Western-style sundials of *Chosun* and traces of projection geometry, Daedong Inst. Korean Stud. 47, 361-396 (2004).
- Kim SH, Mihn BH, Lee MS, Lee YS, A study on the structure innovation of astronomical instruments in *Joseon* Dynasty – focusing on the *Soganui*, the *Ilseongjeongsiui*, and the *Jeokdogyeongwiui*, Publ. Korea Astron. Soc. 27, 61-29 (2012). <https://doi.org/10.5303/PKAS.2012.27.3.061>
- Kim SH, Mihn BH, Lee YS, Analysis on the hour-lines of *Yangcheon-cheok*, J. Astron. Space Sci. 34, 55-65 (2017). <https://doi.org/10.5140/JASS.2017.34.1.55>
- Lee EH, Moon JY, *Gukjo-Yeoksang-Go translated in Korean* (Somyeong Press, Seoul, 2004), 216-230.
- Lee ES, History and fundamental of sundial, Dongbang Hakji 33, 81-156 (1982).
- Lee ES, the horizontal sundial made of marble and crown prince Sohyeon, Korean J. Hist. Sci. Soc. 6, 124-126 (1984).
- Lee ES, The Western-style horizontal sundial made of marble and crown prince Sohyeon, Dongbang Hakji 48, 85-124 (1985).
- Lee GH, Go SW, Choi WH, A technique to improve the readability of ancient inscription by using optical triangulation measurement principle, J. Korea Soc. IT Serv. 11, 103-111 (2012). <https://doi.org/10.9716/KITS.2012.11.sup.103>
- Lee KW, Ahn YS, Mihn BH, Lim YR, Study on the period of the use of *Datong-li* in Korea, J. Astron. Space Sci. 27, 55-68 (2010). <https://doi.org/10.5140/JASS.2010.27.1.055>
- Lee YS, Kim SH, A Study for the restoration of the sundials in King Sejong Era, J. Astron. Space Sci. 28, 143-153 (2011). <https://doi.org/10.5140/JASS.2011.28.2.143>
- Lee YS, Kim SH, Mihn BH, Family of the sun-and-stars time-determining instruments (*Ilseong-jeongsi-ui*) invented during the *Joseon* Dynasty, J. Astron. Space Sci. 33, 237-246 (2016). <https://doi.org/10.5140/JASS.2016.33.3.237>
- Rohr RRJ, Sundial: History, Theory, and Practice (Dover publications, New York, 1996), 49-52.
- Song SY, Yu GN, Park SR, Lee SG, Seong YG, et al., Our science cultural properties (Korea Science and Technology Promotion Foundation, Seoul, 1994), 49-52.
- Yu KL, The first Korea – Japan history of science seminar: on the introduction of Chinese calendars during the *Joseon* Dynasty, Korean J. Hist. Sci. Soc. 4, 98-101 (1982).
- Yu KL, Korean astronomy in the *Joseon* late period, Korean J. Hist. Sci. Soc. 5, 126-131 (1983).